arXiv:0812.4861v2 [math.CO] 8 Jul 2009

On Potentially $K_6 - C_5$ graphic Sequences *

Zhenghua Xu, Chunhui Lai
Department of Mathematics, Zhangzhou Teachers College,
Zhangzhou, Fujian 363000, P. R. of CHINA.
xzh2002528@tom.com(Zhenghua Xu)
zjlaichu@public.zzptt.fj.cn(Chunhui Lai, Corresponding author)

Abstract

For given a graph H, a graphic sequence $\pi = (d_1, d_2, \cdots, d_n)$ is said to be potentially H-graphic if there exists a realization of π containing H as a subgraph. In this paper, we characterize the potentially $K_6 - C_5$ -graphic sequences .

Key words: graph; degree sequence; potentially H-graphic sequences
AMS Subject Classifications: 05C07

1 Introduction

Let G be a simple undirected graph, without loops or multiple edges. Any undefined notation follows that of Bondy and Murty [1]. The set of all non-increasing nonnegative integer sequence $\pi = (d_1, d_2, \cdots, d_n)$ is denoted by NS_n. A sequence π is called graphic if there is a (simple) graph G of order n having degree sequence π and such a graph G is referred as a realization of π. The set of all graphic sequence in NS_n is denoted by GS_n. A graphic sequence π is potentially H-graphic if there is a realization of π containing H as a subgraph. Let C_k and P_k denote a cycle on k vertices and a path on $k+1$ vertices, respectively. Let $\sigma(\pi)$ be the sum of all the

---

*Project Supported by NSF of Fujian(2008J0209), Science and Technology Project of Fujian, Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering" , Project of Fujian Education Department and Project of Zhangzhou Teachers College(SK07009).

terms of π. Let $G - H$ denote the graph obtained from G by removing the edges set $E(H)$ where H is a subgraph of G. In the degree sequence, r^t means that there are t vertices of degree r in the realization of the sequence.

Turán number is the maximum number of edges of a graph with n vertices not containing H as a subgraph and it is denoted by $ex(n, H)$. In terms of graphic sequences, the number $2ex(n, H) + 2$ is the minimum even integer l such that every n-term graphical sequence π with $\sigma(\pi) \geq l$ is forcibly H-graphical. Gould, Jacobson and Lehel [3] considered the following variation of the classical Turán-type extremal problems: determine the smallest even integer $\sigma(H, n)$ such that every n-term positive graphic sequence $\pi = (d_1, d_2, \cdots, d_n)$ with $\sigma(\pi) \geq \sigma(H, n)$ has a realization G containing H as a subgraph. They gave that $\sigma(pK_2, n) = (p - 1)(2n - p) + 2$ for $p \geq 2$; $\sigma(C_4, n) = 2[\frac{3n-1}{2}]$ for $n \geq 4$. Erdös, Jacobson and Lehel [2] showed that $\sigma(K_k, n) \geq (k-2)(2n-k+1)+2$ and conjectured the equality holding and proved the conjecture is true for $k = 3$ and $n \geq 6$. The conjecture is confirmed in [3] and [15]-[18]. Lai[9] determined $\sigma(K_4 - e, n)$ for $n \geq 4$. Yin[25] and Lai[12] determined $\sigma(K_{1,1,3}, n)$ for $n \geq 5$ independently. Yin,Li and Mao[27] determined $\sigma(K_{r+1} - e, n)$ for $r \geq 3$, $r + 1 \leq n \leq 2r$ and $\sigma(K_5 - e, n)$ for $n \geq 5$. Yin and Li[26] gave a good method of determining $\sigma(K_{r+1} - e, n)$ for $r \geq 2$ and $n \geq 3r^2 - r - 1$ (In fact, Yin and Li[26] also determined $\sigma(K_{r+1} - ke, n)$ for $r \geq 2$ and $n \geq 3r^2 - r - 1$). After reading[26], Yin [31] determined $\sigma(K_{r+1} - K_3, n)$ for $n \geq 3r + 5, r \geq 3$.Lai [10-11] determined $\sigma(K_5 - C_4, n), \sigma(K_5 - P_3, n)$ and $\sigma(K_5 - P_4, n)$ where $n \geq 5$. Determining $\sigma(K_{r+1} - H, n)$ where H is a tree on 4 vertices ,is more useful than a cycle on 4 vertices (for example, although $C_4 \not\subset C_i$, $P_3 \subset C_i$ for $i \geq 5$). After reading[26] and [31], Lai and Hu[13] determined $\sigma(K_{r+1} - H, n)$ for $n \geq 4r + 10, r \geq 3, r + 1 \geq k \geq 4$ where H is a graph on k vertices which contains a tree on 4 vertices but not contain a cycle on 3 vertices and $\sigma(K_{r+1} - P_2, n)$ for $n \geq 4r + 8, r \geq 3$. Recently, Lai [14] also determined $\sigma(K_{r+1} - Z_4, n)$, $\sigma(K_{r+1} - (K_4 - e), n)$, $\sigma(K_{r+1} - K_4, n)$ for $n \geq 5r + 16$, $r \geq 4$ and $\sigma(K_{r+1} - Z, n)$ for $n \geq 5r + 19$, $r + 1 \geq k \geq 5$, $j \geq 5$ where Z is a graph on k vertices and j edges which contains a graph Z_4 but not contain a cycle on 4 vertices.

A harder question is to characterize the potentially H-graphic sequences without zero terms. Luo [20] characterized the potentially C_k-graphic sequences for each $k = 3, 4, 5$ and Luo and Warner [21] characterized the potentially K_4-graphic sequences. Lately, Eschen and Niu [22] characterized the potentially $K_4 - e$-graphic sequences. Yin and Chen [29] characterized the potentially $K_{r,s}$-graphic sequences for $r = 2, s = 3$ and $r = 2, s = 4$. Yin et al. [32] characterized the potentially $K_5 - e$ and K_6-graphic sequences. Recently, Hu et al. [4-6] characterized the potentially $K_5 - C_4$, $K_5 - P_4$ and $K_5 - E_3$-graphic sequences where E_3 denotes graphs with 5 vertices and 3 edges and the potentially $K_{3,3}$, $K_6 - C_6$-graphic sequences in [7]. Yin [33] characterized the potentially $K_6 - K_3$-graphic sequences. In this paper, we have characterized the potentially $K_6 - C_5$ -graphic sequences.

2 Preparations

Let $\pi = (d_1, \cdots, d_n) \epsilon NS_n, 1 \leq k \leq n$. Let

$$\pi''_k = \begin{cases} (d_1 - 1, \cdots, d_{k-1} - 1, d_{k+1} - 1, \cdots, d_{d_k+1} - 1, d_{d_k+2}, \cdots, d_n), \\ \text{ if } d_k \geq k, \\ (d_1 - 1, \cdots, d_{d_k} - 1, d_{d_k+1}, \cdots, d_{k-1}, d_{k+1}, \cdots, d_n), \\ \text{if } d_k < k. \end{cases}$$

Denote $\pi'_k = (d'_1, d'_2, \cdots, d'_{n-1})$, where $d'_1 \geq d'_2 \geq \cdots \geq d'_{n-1}$ is a rearrangement of the $n - 1$ terms of π''_k. Then π'_k is called the residual sequence obtained by laying off d_k from π. For simplicity, we denote π'_n by π' in this paper.

If $S \subseteq NS_n$, then $\pi \in S$ if and only if π is an element of S by rearranging the terms of π.

Let $\pi_1 = (d^1_1, d^1_2, \cdots, d^1_n) \in NS_n$, $\pi_2 = (d^2_1, d^2_2, \cdots, d^2_m) \in NS_n$ and $n \geq m$, $d^1_i \geq d^2_i (i = 1, 2, \cdots, m)$. Let $\pi^* = \pi_1 - \pi_2 = (d^1_1 - d^2_1, d^1_2 - d^2_2, \cdots, d^1_m - d^2_m, d^1_{m+1}, \cdots, d^1_n)$.

For a non-increasing positive integer sequence $\pi = (d_1, d_2, \cdots, d_n)$, we write $m(\pi)$ and $h(\pi)$ to denote the largest positive terms of π and the smallest positive terms of π, respectively. We need the following the results.

Theorem 2.1 [3] If $\pi = (d_1, d_2, \cdots, d_n)$ is a graphic sequence with a realization G containing H as a subgraph, then there exists a realization G' of π containing H as a subgraph so that the vertices of H have the largest degrees of π.

Theorem 2.2 [19] If $\pi = (d_1, d_2, \cdots, d_n)$ is a sequence of nonnegative integers where $1 \leq m(\pi) \leq 2$, $h(\pi) = 1$ and $\sigma(\pi)$ is even, then π is graphic.

Lemma 2.3 (Kleitman and Wang [8]) π is graphic if and only if π' is graphic.

The following corollary is obvious.

Corollary 2.4 Let H be a simple graph. If π' is potentially H-graphic, then π is potentially H-graphic.

Lemma 2.5 [24] Let G be a simple undirected graph with the vertex v_i and $d_G(v'_i) = d'_i(i = 1, 2, \cdots, m)$, $d'_1 \geq d'_2 \geq \cdots \geq d'_m$. If $\pi = (d_1, d_2, \cdots, d_n) \in NS_n$ is potentially G-graphic and $n-m \geq (d_1 - d'_m)$, then there exists a realization H of π containing G as a subgraph so that the vertices of G have the largest degrees of π in proper order, that is, if the vertices of G are v'_i, $i = 1, 2, \cdots, m(m \leq n)$, $d_G(v'_i) = d'_i$ and $d'_1 \geq d'_2 \geq \cdots \geq d'_m$, then there exists a realization H of π containing G as a subgraph so that the vertices of H are v_i, $i = 1, 2, \cdots, n$, $d_H(v_i) = d_i$, $G \subseteq H[v_1, v_2, \cdots, v_m]$ and the vertices v'_i are corresponding to the vertices v_i, $i = 1, 2, \cdots, m$.

Lemma 2.6 [23] Let $\pi = (4^{x_1}, 3^{x_2}, 2^{x_3}, 1^{x_4})$ where $\sigma(\pi)$ is even, $x_1 + x_2 + x_3 + x_4 = n$ and $n \geq 1$. Then $\pi \in GS_n$ if and only if $\pi \notin S$, where $S = \{(2),(2^2),(3,1),(3^2),(3,2,1),(3^2,2),(3^3,1),(3^2,1^2),(4),(4,1^2),(4,2),(4,2^2),(4,2^3),(4,2,1^2),(4,3^2),(4,3^2,2),(4,3,1),(4,3,1^3),(4,3^2,1^2),(4,3,2,1),(4^2),(4^2,1^2),(4^2,1^4),(4^2,2,1^2),(4^2,2),(4^2,2^2),(4^2,3^2),(4^2,3,1),(4^2,3,1^3),(4^2,3,2,1),(4^3),(4^3,1^2),(4^3,2,1^2),(4^3,1^4),(4^3,2),(4^3,2^2),(4^3,3,1),(4^4),(4^4,1^2),(4^4,2)\}$.

In order to prove Theorem 3.1, at first, we prove Lemma 2.7 to Lemma 2.11. Let $\pi_1^1 = (5, 3^5)$ be a graphic sequence of $K_6 - C_5$ and $\pi^* = \pi - \pi_1^1$.

Lemma 2.7 (1) If $\pi = (5^4, 4^{n-4})$ where $n \geq 6$ and $\sigma(\pi)$ is even, then π is potentially $K_6 - C_5$-graphic;

(2) If $\pi = (5^3, 4^i, 3^j, 2^{n-3-i-j})$ where $n \geq 6$, $i + j \geq 3$ and $\sigma(\pi)$ is even, then π is potentially $K_6 - C_5$-graphic if and only if $\pi \neq (5^3, 3^3)$.

Proof: (1) If $\pi = (5^4, 4^{n-4})$ where $n \geq 6$, then $\pi^* = (2, 2, 2, 1, 1, 4^{n-6})$. Let $\pi_1 = (4^{n-6})$. If $\pi_1 = (4^{n-6}) \notin S$, according to Lemma 2.6, then π_1 is graphic. Let G be a realization of $\pi_1 = (4^{n-6})$, then $(K_6 - e) \bigcup G$ is a realization of $\pi = (5^4, 4^{n-4})$, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (4^{n-6}) \in S$, then $\pi_1 =(4)$, (4^2), (4^3), (4^4). Therefore, $\pi =(5^4, 4^3)$, $(5^4, 4^4)$, $(5^4, 4^5)$, $(5^4, 4^6)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic.

(2) If $\pi = (5^3, 4^i, 3^j, 2^{n-3-i-j})$ where $n \geq 6$ and $i + j \geq 3$, then we consider the following cases.

Case 1: $i = 0$

If $\pi = (5^3, 3^j, 2^{n-3-j})(j \geq 3)$, then $\pi^* = (2^2, 3^{j-3}, 2^{n-3-j})$.

If j=3, then $\pi^* = (2^2, 2^{n-6})$. If $n = 6$, then $\pi = (5^3, 3^3)$. It is easy to check that π =$(5^3, 3^3)$ is not potentially $K_6 - C_5$-graphic. If $n \geq 7$, then let $\pi_1 = (1, 1, 2^{n-7})$. Since $\pi_1 = (1, 1, 2^{n-7}) \notin S$, according to Lemma 2.6, then π_1 is graphic. Firstly, let G_1 be a realization of $\pi_1 = (1, 1, 2^{n-7})$ and u_1 and u_2 denote two vertices of degree 1 in G_1. Secondly, we add a vertex u_3 to G_1. Let u_3 be adjacent to u_1 and u_2. Then we obtain a graph G_2 and G_2 is a realization of $\pi^* = (2^2, 2^{n-6})(n \geq 7)$. Finally, let v_1 and v_2 denote two vertices of degree 3 and be not adjacent in $K_6 - C_5$. We replace v_1 and u_1 with a single vertex whose incident edges are the edges that were incident to v_1 or u_1 and we also replace v_2 and u_2 with a single vertex whose incident edges are the edges that were incident to v_2 or u_2 in $G_2 \bigcup (K_6 - C_5)$. Then we obtain a graph G and G is a realization of $\pi = (5^3, 3^3, 2^{n-6})$, hence π is potentially $K_6 - C_5$-graphic. We similarly do as follows.

If $j \geq 4$, then let $\pi_1 = (1, 2, 3^{j-4}, 2^{n-3-j})$. If $\pi_1 = (1, 2, 3^{j-4}, 2^{n-3-j}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (1, 2, 3^{j-4}, 2^{n-3-j}) \in S$, then π_1 =$(3, 2, 1)$. Therefore, π =$(5^3, 3^5)$. It is easy to check that π =$(5^3, 3^5)$ is potentially $K_6 - C_5$-graphic .

Case 2: $i = 1$

If $\pi = (5^3, 4, 3^j, 2^{n-4-j})(j \geq 2)$, then $\pi^* = (2, 2, 1, 3^{j-2}, 2^{n-4-j})$. Let $\pi_1 = (1, 3^{j-2}, 2^{n-4-j})$. If $\pi_1 = (1, 3^{j-2}, 2^{n-4-j}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (1, 3^{j-2}, 2^{n-4-j}) \in S$, then π_1 =$(3, 1)$, $(3, 2, 1)$, $(3^3, 1)$. Therefore, π =$(5^3, 4, 3^3)$,$(5^3, 4, 3^3, 2)$,$(5^3, 4, 3^5)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic .

Case 3: $i = 2$

If $\pi = (5^3, 4^2, 3^j, 2^{n-5-j})(j \geq 1)$, then $\pi^* = (2, 2, 1, 1, 3^{j-1}, 2^{n-5-j})$. Let $\pi_1 = (3^{j-1}, 2^{n-5-j})$. If $\pi_1 = (3^{j-1}, 2^{n-5-j}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (3^{j-1}, 2^{n-5-j}) \in S$, then π_1 =(2), (2^2), (3^2), $(3^2, 2)$. Therefore, $\pi = (5^3, 4^2, 3, 2)$, $(5^3, 4^2, 3, 2^2)$, $(5^3, 4^2, 3^3)$, $(5^3, 4^2, 3^3, 2)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic .

Case 4: $i \geq 3$

If $\pi = (5^3, 4^i, 3^j, 2^{n-3-i-j})$ $(j \geq 0)$, then $\pi^* = (2, 2, 1, 1, 1, 4^{i-3}, 3^j, 2^{n-3-i-j})$. Let $\pi_1 = (1, 4^{i-3}, 3^j, 2^{n-3-i-j})$. If $\pi_1 = (1, 4^{i-3}, 3^j, 2^{n-3-i-j}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (1, 4^{i-3}, 3^j, 2^{n-3-i-j}) \in S$, then π_1 =$(3, 1)$,

$(3,2,1)$, $(3^3,1)$, $(4,3,1)$, $(4,3,2,1)$, $(4^2,3,1)$, $(4^2,3,2,1)$, $(4^3,3,1)$. Therefore, $\pi =(5^3,4^3,3)$, $(5^3,4^3,3,2)$,$(5^3,4^3,3^3)$,$(5^3,4^4,3)$,$(5^3,4^4,3,2)$, $(5^3,4^5,3)$, $(5^3,4^5,3,2)$, $(5^3,4^6,3)$. It is easy to check that all of these are potentially K_6-C_5.

Lemma 2.8 If $\pi=(5^2,4^i,3^j,2^{n-2-i-j})$ where $i+j\geq 4$, $n\geq 6$ and $\sigma(\pi)$ is even, then π is potentially K_6-C_5-graphic if and only if $\pi \neq(5^2,3^4)$, $(5^2,3^4,2)$,$(5^2,3^6)$,$(5^2,4,3^4)$.

Proof: If $\pi=(5^2,4^i,3^j,2^{n-2-i-j})$ where $i+j\geq 4$ and $n\geq 6$, then we consider the following cases.

Case 1: $i=0$

If $\pi=(5^2,3^j,2^{n-2-j})(j\geq 4)$, then $\pi^*=(2,3^{j-4},2^{n-2-j})$. If $\pi^*\notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially K_6-C_5-graphic. If $\pi^*\in S$, then $\pi^*=(2)$,(2^2),$(3^2,2)$. Therefore, $\pi=(5^2,3^4)$,$(5^2,3^4,2)$,$(5^2,3^6)$. It is easy to check that these are not potentially K_6-C_5-graphic .

Case 2: $i=1$

If $\pi=(5^2,4,3^j,2^{n-3-j})(j\geq 3)$, then $\pi^*=(2,1,3^{j-3},2^{n-3-j})$. Let $\pi_1=(1,3^{j-3},2^{n-3-j})$. If $\pi_1=(1,3^{j-3},2^{n-3-j})\notin S$, according to Lemma 2.6,then π_1 is graphic, hence π is potentially K_6-C_5-graphic. If $\pi_1=(1,3^{j-3},2^{n-3-j})\in S$,then $\pi_1=(3,1)$,$(3,2,1)$,$(3^3,1)$.Therefore, $\pi=(5^2,4,3^4)$, $(5^2,4,3^4,2)$,$(5^2,4,3^6)$. It is easy to check that others are potentially K_6-C_5-graphic except for $(5^2,4,3^4)$.

Case 3: $i=2$

If $\pi=(5^2,4^2,3^j,2^{n-4-j})(j\geq 2)$, then $\pi^*=(2,1,1,3^{j-2},2^{n-4-j})$. Let $\pi_1=(3^{j-2},2^{n-4-j})$. If $\pi_1=(3^{j-2},2^{n-4-j})\notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially K_6-C_5-graphic. If $\pi_1=(3^{j-2},2^{n-4-j})\in S$,then $\pi_1=(2)$,(2^2),(3^2),$(3^2,2)$.Therefore,$\pi=(5^2,4^2,3^2,2)$, $(5^2,4^2,3^2,2^2)$, $(5^2,4^2,3^4)$, $(5^2,4^2,3^4,2)$. It is easy to check that all of these are potentially K_6-C_5-graphic .

Case 4: $i=3$

If $\pi=(5^2,4^3,3^j,2^{n-5-j})(j\geq 1)$, then $\pi^*=(2,1,1,1,3^{j-1},2^{n-5-j})$. Let $\pi_1=(1,3^{j-1},2^{n-5-j})$. If $\pi_1=(1,3^{j-1},2^{n-5-j})\notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially K_6-C_5-graphic. If $\pi_1=(1,3^{j-1},2^{n-5-j})\in S$, then $\pi_1=(3,1)$, $(3,2,1)$, $(3^3,1)$. Therefore, $\pi=(5^2,4^3,3^2)$, $(5^2,4^3,3^2,2)$,$(5^2,4^3,3^4)$. It is easy to check that all of these are potentially K_6-C_5-graphic .

Case 5: $i\geq 4$

If $\pi = (5^2, 4^i, 3^j, 2^{n-2-i-j})$ $(j \geq 0)$, then $\pi^* = (2, 1, 1, 1, 1, 4^{i-4}, 3^j, 2^{n-2-i-j})$. Let $\pi_1 = (4^{i-4}, 3^j, 2^{n-2-i-j})$. If $\pi_1 = (4^{i-4}, 3^j, 2^{n-2-i-j}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (4^{i-4}, 3^j, 2^{n-2-i-j}) \in S$, then $\pi_1 =(2)$, (2^2), (3^2), $(3^2, 2)$, (4), $(4, 2)$, $(4, 2^2)$, $(4, 2^3)$, $(4, 3^2)$, $(4, 3^2, 2)$, (4^2), $(4^2, 2)$, $(4^2, 2^2)$, $(4^2, 3^2)$, (4^3), $(4^3, 2)$, $(4^3, 2^2)$, (4^4), $(4^4, 2)$. Therefore, $\pi =(5^2, 4^4, 2)$, $(5^2, 4^4, 2^2)$, $(5^2, 4^4, 3^2)$, $(5^2, 4^4, 3^2, 2)$, $(5^2, 4^5)$, $(5^2, 4^5, 2)$, $(5^2, 4^5, 2^2)$, $(5^2, 4^5, 2^3)$, $(5^2, 4^5, 3^2)$, $(5^2, 4^5, 3^2, 2)$, $(5^2, 4^6)$, $(5^2, 4^6, 2)$, $(5^2, 4^6, 2^2)$, $(5^2, 4^6, 3^2)$, $(5^2, 4^7)$, $(5^2, 4^7, 2)$, $(5^2, 4^7, 2^2)$, $(5^2, 4^8)$, $(5^2, 4^8, 2)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic .

Lemma 2.9 If $\pi = (5, 4^i, 3^j, 2^k, 1^{n-i-j-k-1})$ where $i + j \geq 5$, $n \geq 6$ and $\sigma(\pi)$ is even, then π is potentially $K_6 - C_5$-graphic if and only if $\pi \neq (5, 3^5, 2)$, $(5, 3^5, 2^2)$, $(5, 3^7)$, $(5, 3^6, 1)$, $(5, 3^6, 2, 1)$, $(5, 3^7, 2)$, $(5, 3^8, 1)$, $(5, 3^7, 1^2)$, $(5, 4, 3^5)$, $(5, 4, 3^5, 2)$, $(5, 4, 3^7)$, $(5, 4, 3^6, 1)$, $(5, 4^2, 3^5)$.

Proof: If $\pi = (5, 4^i, 3^j, 2^k, 1^{n-i-j-k-1})$ where $i + j \geq 5$ and $n \geq 6$, then we consider the following cases.

Case 1: $i = 0$

If $\pi = (5, 3^j, 2^k, 1^{n-j-k-1})(j \geq 5)$, then $\pi^* = (3^{j-5}, 2^k, 1^{n-j-k-1})$. If $\pi^* \notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi^* \in S$, then $\pi^* =(2)$, (2^2), (3^2), $(3, 1)$, $(3, 2, 1)$,$(3^2, 2)$,$(3^3, 1)$,$(3^2, 1^2)$. Therefore, $\pi =(5, 3^5, 2)$, $(5, 3^5, 2^2)$, $(5, 3^7)$, $(5, 3^6, 1)$, $(5, 3^6, 2, 1)$, $(5, 3^7, 2)$, $(5, 3^8, 1)$, $(5, 3^7, 1^2)$. It is easy to check that these are not potentially $K_6 - C_5$-graphic.

Case 2: $i = 1$

If $\pi = (5, 4, 3^j, 2^k, 1^{n-2-j-k})(j \geq 4)$, then $\pi^* = (1, 3^{j-4}, 2^k, 1^{n-2-j-k})$. If $\pi^* \notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially K_6-C_5-graphic. If $\pi^* = (1, 3^{j-4}, 2^k, 1^{n-2-j-k}) \in S$, then $\pi^* =(3, 1)$,$(3, 2, 1)$, $(3^3, 1)$,$(3^2, 1^2)$. Therefore,$\pi =(5, 4, 3^5)$,$(5, 4, 3^5, 2)$, $(5, 4, 3^7)$,$(5, 4, 3^6, 1)$. It is easy to check that these are not potentially $K_6 - C_5$-graphic.

Case 3: $i = 2$

If $\pi = (5, 4^2, 3^j, 2^k, 1^{n-3-j-k})(j \geq 3)$,then $\pi^* = (1, 1, 3^{j-3}, 2^k, 1^{n-3-j-k})$. Let $\pi_1 = (3^{j-3}, 2^k, 1^{n-3-j-k})$. If $\pi_1 = (3^{j-3}, 2^k, 1^{n-3-j-k}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (3^{j-3}, 2^k, 1^{n-3-j-k}) \in S$, then $\pi_1 =(2)$, (2^2), (3^2), $(3, 1)$, $(3, 2, 1)$, $(3^2, 2)$, $(3^3, 1)$, $(3^2, 1^2)$. Therefore, $\pi =(5, 4^2, 3^3, 2)$, $(5, 4^2, 3^3, 2^2)$, $(5, 4^2, 3^5)$, $(5, 4^2, 3^4, 1)$, $(5, 4^2, 3^4, 2, 1)$, $(5, 4^2, 3^5, 2)$,$(5, 4^2, 3^6, 1)$,$(5, 4^2, 3^5, 1^2)$. It is easy to check that others are potentially $K_6 - C_5$-graphic except for $(5, 4^2, 3^5)$.

Case 4: $i = 3$

If $\pi = (5, 4^3, 3^j, 2^k, 1^{n-4-j-k})$ $(j \geq 2)$, then $\pi^* = (1, 1, 1, 3^{j-2}, 2^k, 1^{n-4-j-k})$. Let $\pi_1 = (1, 3^{j-2}, 2^k, 1^{n-4-j-k})$. If $\pi_1 = (1, 3^{j-2}, 2^k, 1^{n-4-j-k}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (1, 3^{j-2}, 2^k, 1^{n-4-j-k}) \in S$, then $\pi_1 =(3, 1)$, $(3, 2, 1)$, $(3^3, 1)$, $(3^2, 1^2)$. Therefore, $\pi =(5, 4^3, 3^3)$, $(5, 4^3, 3^3, 2)$, $(5, 4^3, 3^5)$,$(5, 4^3, 3^4, 1)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic .

Case 5: $i = 4$

If $\pi = (5, 4^4, 3^j, 2^k, 1^{n-5-j-k})$ $(j \geq 1)$, then $\pi^* = (1, 1, 1, 1, 3^{j-1}, 2^k, 1^{n-5-j-k})$. Let $\pi_1 = (3^{j-1}, 2^k, 1^{n-5-j-k})$. If $\pi_1 = (3^{j-1}, 2^k, 1^{n-5-j-k}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (3^{j-1}, 2^k, 1^{n-5-j-k}) \in S$, then $\pi_1 =(2)$, (2^2), (3^2), $(3, 1)$, $(3, 2, 1)$, $(3^2, 2)$, $(3^3, 1)$, $(3^2, 1^2)$. Therefore, $\pi =(5, 4^4, 3, 2)$, $(5, 4^4, 3, 2^2)$, $(5, 4^4, 3^3)$, $(5, 4^4, 3^2, 1)$, $(5, 4^4, 3^2, 2, 1)$, $(5, 4^4, 3^3, 2)$, $(5, 4^4, 3^4, 1)$, $(5, 4^4, 3^3, 1^2)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic .

Case 6: $i \geq 5$

If $\pi = (5, 4^i, 3^j, 2^k, 2^{n-1-i-j-k})$($j \geq 0$ and $\sigma(\pi)$ is even), then π is a graphic sequence, hence $\pi^* = (1, 1, 1, 1, 1, 4^{i-5}, 3^j, 2^k, 1^{n-1-i-j-k})$. Let $\pi_1 = (1, 4^{i-5}, 3^j, 2^k, 1^{n-1-i-j-k})$. If $\pi_1 = (1, 4^{i-5}, 3^j, 2^k, 1^{n-1-i-j-k}) \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 = (1, 4^{i-5}, 3^j, 2^k, 1^{n-1-i-j-k}) \in S$, then $\pi_1 =(3, 1)$, $(3, 2, 1)$, $(3^3, 1)$, $(3^2, 1^2)$, $(4, 1^2)$, $(4, 2, 1^2)$, $(4, 3, 1)$, $(4, 3, 1^3)$, $(4, 3^2, 1^2)$, $(4, 3, 2, 1)$,$(4^2, 1^2)$,$(4^2, 1^4)$,$(4^2, 2, 1^2)$, $(4^2, 3, 1)$,$(4^2, 3, 1^3)$,$(4^2, 3, 2, 1)$,$(4^3, 1^2)$, $(4^3, 2, 1^2)$, $(4^3, 1^4)$, $(4^3, 3, 1)$, $(4^4, 1^2)$. Therefore, $\pi =(5, 4^5, 3)$, $(5, 4^5, 3, 2)$, $(5, 4^5, 3^3)$, $(5, 4^5, 3^2, 1)$, $(5, 4^6, 1)$, $(5, 4^6, 2, 1)$, $(5, 4^6, 3)$, $(5, 4^6, 3, 1^2)$, $(5, 4^6, 3^2, 1)$, $(5, 4^6, 3, 2)$, $(5, 4^7, 1)$, $(5, 4^7, 1^3)$, $(5, 4^7, 2, 1)$, $(5, 4^7, 3)$, $(5, 4^7, 3, 1^2)$, $(5, 4^7, 3, 2)$, $(5, 4^8, 1)$, $(5, 4^8, 2, 1)$, $(5, 4^8, 1^3)$, $(5, 4^8, 3)$, $(5, 4^9, 1)$. It is easy to check that all of these are potentially $K_6 - C_5$-graphic .

Lemma 2.10 If $\pi = (d_1, d_2, 3^{n-2})$ where $d_1 \geq 5$, $d_2 \geq 3$, $n \geq 6$ and $\sigma(\pi)$ is even, then

(1) $\pi \in GS_n$;

(2) π is potentially $K_6 - C_5$-graphic if the following conditions hold:

(i) If $n \geq 7$ and n is even, then $\pi \neq ((n-1)^2, 3^{n-2})$ and $\pi \neq ((n-2)^2, 3^{n-2})$;

(ii) If $n \geq 7$ and n is odd, then $\pi \neq ((n-1), (n-2), 3^{n-2})$;

(iii) $\pi \neq (5, 3^7)$,$(5, 4, 3^5)$,$(5, 4, 3^7)$,$(5^2, 3^4)$,$(5^2, 3^6)$, $(6, 3^6)$,$(6, 3^8)$,$(7, 3^7)$,

$(6, 4, 3^6)$.

Proof: (1) is obvious.

(2) If $\pi = (d_1, d_2, 3^{n-2})$ where $d_1 \geq 5$, $d_2 \geq 3$, $n \geq 6$ and $\sigma(\pi)$ is even, then π is potentially $K_6 - C_5$-graphic when $\pi^* = (d_1 - 5, d_2 - 3, 3^{n-6})$ is graphic and two vertices of degree $d_1 - 5$ and $d_2 - 3$ are not adjacent. We consider the following cases.

Case 1: $d_1 - 5 = 0$

If $\pi = (5, d_2, 3^{n-2})$, then $\pi^* = (d_2 - 3, 3^{n-6})$.

If $d_2 - 3 = 0$, then $\pi^* = (3^{n-6})$. If $\pi^* \notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi^* \in S$, then π^* =(3^2). Therefore, $\pi = (5, 3^7)$. It is easy to check that $\pi = (5, 3^7)$ is not potentially $K_6 - C_5$-graphic.

If $d_2 - 3 = 1$, then $\pi^* = (1, 3^{n-6})$. If $\pi^* \notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi^* \in S$,then π^* =$(3, 1)$,$(3^3, 1)$. Therefore, $\pi = (5, 4, 3^5)$,$(5, 4, 3^7)$. It is easy to check that $\pi = (5, 4, 3^5)$ and $(5, 4, 3^7)$ are not potentially $K_6 - C_5$-graphic.

If $d_2 - 3 = 2$, then $\pi^* = (2, 3^{n-6})$. If $\pi^* \notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi^* \in S$, then π^* =(2),$(3^2, 2)$. Therefore, $\pi = (5^2, 3^4)$,$(5^2, 3^6)$. It is easy to check that $\pi = (5^2, 3^4)$ and $(5^2, 3^6)$ are not potentially $K_6 - C_5$-graphic.

Case 2: $d_1 - 5 \neq 0$

If $\pi = (d_1, d_2, 3^{n-2})$, then $\pi^* = (d_1 - 5, d_2 - 3, 3^{n-6})$.

If $d_2 - 3 = 0$, then $\pi^* = (d_1 - 5, 3^{n-6})$. Let π_1 =$(3^{(n-6)-(d_1-5)}, 2^{d_1-5})$. If $\pi_1 \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic. If $\pi_1 \in S$, then π_1 =(2),(2^2),(3^2),$(3^2, 2)$. Therefore, π =$(6, 3^6)$,$(7, 3^7)$,$(5, 3^7)$,$(6, 3^8)$. It is easy to check that π =$(6, 3^6)$,$(7, 3^7)$ and $(6, 3^8)$ are not potentially $K_6 - C_5$-graphic and $\pi = (5, 3^7)$, a contradiction to $d_1 - 5 \neq 0$.

If $1 \leq d_2 - 3 \leq (n - 6) - (d_1 - 5)$, then let $\pi_1 = (3^{(n-6)-(d_1-5)-(d_2-3)}, 2^{(d_1-5)+(d_2-3)})$. If $\pi_1 \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially $K_6 - C_5$-graphic.If $\pi_1 \in S$,then π_1 =(2),(2^2),(3^2),$(3^2, 2)$. Therefore, $\pi = (5, 4, 3^5)$,$(6, 4, 3^6)$,$(5^2, 3^6)$,$(5, 4, 3^7)$. It is easy to check that $\pi = (6, 4, 3^6)$ is not potentially $K_6 - C_5$-graphic and $\pi = (5, 4, 3^5)$,$(5^2, 3^6)$, $(5, 4, 3^7)$, a contradiction to $d_1 - 5 \neq 0$.

If $n - 6 \geq d_2 - 3 > (n - 6) - (d_1 - 5)$, then let $\pi_1 = (2^{(n-6)-(d_1-5)}, 1^{(d_2-3)+(d_1-5)-(n-6)}, 2^{(d_1-5)-(d_2-3)-(d_1-5)+(n-6)}) = (2^{2(n-6)-(d_2-3)-(d_1-5)}, 1^{(d_1-5)+(d_2-3)-(n-6)})$. If $\pi_1 \notin S$, according to Lemma 2.6, then π_1 is graphic, hence π is potentially K_6-C_5-graphic. If $\pi_1 \in S$, then π_1 =(2),(2^2). Therefore, $\pi = (5,4,3^5)$,$(6,4,3^6)$,$(5^2,3^6)$, a contradiction.

If $d_2-3>(n-6)$ and $d_2 \leq n-1$, then $d_2=n-1$ or $d_2=n-2$. If $d_2=n-1$, then $\pi=((n-1)^2,3^{n-2})(n\ is\ even)$. It is easy to check that $\pi = ((n-1)^2,3^{n-2})(n\ is\ even)$ is not potentially K_6-C_5-graphic. If $d_2=n-2$, then $\pi=(n-1,n-2,3^{n-2})(n\ is\ odd)$ or $\pi=(n-2,n-2,3^{n-2})(n\ is\ even)$. It is easy to check that these are not potentially K_6-C_5-graphic.

Lemma 2.11 If $\pi=(d_1,3^5,2^{n-6})$ where $d_1 \geq 5$, $n \geq 6$ and $\sigma(\pi)$ is even, then

(1) $\pi \in GS_n$;

(2) π is potentially K_6-C_5-graphic if and only if $\pi \neq$$(5,3^5,2)$,$(5,3^5,2^2)$.

Proof: (1) is obvious

(2) If $\pi=(d_1,3^5,2^{n-6})$ where $d_1 \geq 5$, $n \geq 6$ and $\sigma(\pi)$ is even, then π is potentially K_6-C_5-graphic if and only if $\pi^*=(d_1-5,2^{n-6})$ is graphic. We consider the following cases.

Case 1: $d_1-5=0$

If $\pi=(5,3^5,2^{n-6})$, then $\pi^*=(2^{n-6})$. If $\pi^* \notin S$, according to Lemma 2.6, then π^* is graphic, hence π is potentially K_6-C_5-graphic. If $\pi^* \in S$, then π^* =(2),(2^2). Therefore, $\pi=(5,3^5,2)$,$(5,3^5,2^2)$. It is easy to check that these are not potentially K_6-C_5-graphic.

Case 2: $1 \leq d_1-5 \leq n-6$

If $\pi=(d_1,3^5,2^{n-6})$, then let $\pi_1=(2^{(n-6)-(d_1-5)},1^{d_1-5})$. Since $\pi_1 \notin S$, according to Lemma 2.6, π_1 is graphic, hence π is potentially K_6-C_5-graphic.

3 Main Theorems

Theorem 3.1 Let $\pi=(d_1,d_2,\cdots,d_n) \in GS_n$ and $n \geq 6$. Then π is potentially K_6-C_5-graphic if and only if the following conditions hold:

(1) $d_1 \geq 5, d_6 \geq 3$;

(2) If $\pi=(d_1,d_2,d_3,3^i,2^j,1^{n-i-j-3})$ where $i \geq 3$ and $d_3 \geq 5$, then $d_1+d_2+d_3 \leq n+2i+j+1$;

(3) If $\pi=(d_1,d_2,3^4,2^j,1^{n-j-6})$ where $j \geq 0$ and $d_2 \geq 5$, then $d_1+d_2 \leq n+j+2$;

(4) If $\pi=(d_1,d_2,d_3,3^i,2^j,1^{n-i-j-3})$ where $n \geq 8$, $i \geq 4$ and $d_3 \geq 4$, then

(i) if $n=i+3$, $d_1=n-1$ and $d_3 \geq 5$, then $d_2 \leq n-2$;

(ii) if $n \geq i+j+4$, $d_1 \geq i+j+3$, $d_2 \geq max\{d_3+2, i+j+2\}$ and $d_3 = 4$, then $d_1 + d_2 \leq n+i+j$;

(iii) if $n \geq i+j+4$, $d_1 = n-1$, $d_2 \geq d_3+2$ and $d_3 \geq 5$, then $d_1 + d_2 \leq n+i+j$;

(iv) if $n = i+j+3$, $j \geq 1$, $d_1 = n-1$ and $d_2 \geq d_3+2$, then $d_2 \leq n-2$;

(5) If $\pi = (d_1, d_2, 3^i, 2^j, 1^{n-i-j-2})$ where $i \geq 5$ and $d_2 \geq 5$, then

(i) if $n \geq i+1$, $j \geq 2$ or $j = 0$, $d_1 \geq i+j$, then $d_1 + d_2 \leq n+i+j-2$;

(ii) if $d_1 = i+j+1$ and $d_1 \geq n-2$, then $d_2 \leq n-3$;

(iii) if $d_1 = i+j = n-2$ and $i \geq 6$, then $d_2 \leq n-3$;

(6) $\pi \neq ((n-1)^2, 4, 3^{n-3})(n \geq 7 \ \ and \ \ \ n \ \ is \ \ odd)$;

(7) $\pi \neq (n-1, 3^6, 1^{n-7})(n \geq 7)$,$(n-1, 3^7, 1^{n-8})(n \geq 8)$,$(5, 4, 3^5)$,$(5^2, 3^5, 1)$, $(5, 4^2, 3^5)$, $(5, 4, 3^7)$, $(6, 3^7, 1)$, $(5^2, 3^6)$, $(5^2, 4, 3^4)$, $(5, 3^7)$, $(6, 4, 3^6)$, $(5^2, 3^4, 2)$, $(5, 3^6, 1)$, $(5, 3^6, 2, 1)$, $(5, 3^7, 2)$, $(5, 3^8, 1)$, $(5, 4, 3^5, 2)$, $(5, 4, 3^6, 1)$, $(5, 3^5, 2)$, $(5, 3^5, 2^2)$, $(6, 5, 3^5)$, $(5, 3^7, 1^2)$, $(6, 3^8)$, $(6^2, 3^4, 2)$.

Proof: First we show the conditions (1) to (7) are necessary conditions for π to be potentially $K_6 - C_5$-graphic. Assume that π is potentially $K_6 - C_5$-graphic.

(1) is obvious.

(2) If $\pi = (d_1, d_2, d_3, 3^i, 2^j, 1^{n-i-j-3})$ where $i \geq 3$ and $d_3 \geq 5$, then $(d_1 + d_2 + d_3 - 11) - 2 \leq 3 \times (i-3) + 2 \times j + 1 \times (n-i-j-3)$, i.e., $d_1 + d_2 + d_3 \leq n + 2i + j + 1$.

(3) If $\pi = (d_1, d_2, 3^4, 2^j, 1^{n-j-6})$ where $j \geq 0$ and $d_2 \geq 5$, then $(d_1 + d_2 - 8) \leq 2 \times j + (n-j-6)$, i.e., $d_1 + d_2 \leq n + j + 2$.

(4) If $\pi = (d_1, d_2, d_3, 3^i, 2^j, 1^{n-i-j-3})$ where $n \geq 8$, $i \geq 4$ and $d_3 \geq 4$, then

(i) if $n = i+3$, $d_1 = n-1$ and $d_3 \geq 5$, then $d_2 - 3 - 1 \leq n-6$, i.e., $d_2 \leq n-2$;

(ii) if $n \geq i+j+4$, $d_1 \geq i+j+3$, $d_2 \geq max\{d_3+2, i+j+2\}$ and $d_3 = 4$, then $(d_2 - 3) - 1 \leq (i-3) + j + (n-i-j-3) + (i-3) + j - (d_1 - 5)$ or $(d_1 - 3) - 1 \leq (i-3) + j + (n-i-j-3) + (i-3) + j - (d_2 - 5)$, i.e., $d_1 + d_2 \leq n+i+j$.

(iii) if $n \geq i+j+4$, $d_1 = n-1$, $d_2 \geq d_3+2$ and $d_3 \geq 5$, then $(d_2 - 3) - 1 \leq (i-3) + j + (n-i-j-3) + (i-3) + j - (d_1 - 5)$, i.e., $d_1 + d_2 \leq n+i+j$ or $(d_1 - 3) - 1 \leq (i-3) + j + (n-i-j-3) + (i-3) + j - (d_2 - 5)$, i.e., $d_1 + d_2 \leq n+i+j$;

(iv) if $n = i+j+3$, $j \geq 1$, $d_1 = n-1$, $d_2 \geq d_3+2$, then $d_2 - 3 - 1 \leq (i-3) + j + (n-i-j-3) = n-6$, i.e., $d_2 \leq n-2$.

(5) If $\pi = (d_1, d_2, 3^i, 2^j, 1^{n-i-j-2})$ where $i \geq 5$ and $d_2 \geq 5$, then

(i) if $n \geq i+1$, $j \geq 2$ or $j = 0$, $d_1 \geq i+j$, then $d_2 - 3 \leq (i-4) + j + (n-i-j-2) - [(d_1-5) - (i-4) - j]$ or $d_1 - 5 \leq (i-4) + j + (n-i-j-2) - [(d_2-3) - (i-4) - j]$, i.e., $d_1 + d_2 \leq n + i + j - 2$;

(ii) if $d_1 = i+j+1$ and $d_1 \geq n-2$, then $\pi^* = (d_1-5, d_2-3, 3^{i-4}, 2^j, 1^{n-i-j-2})$. Therefore, $d_2 - 3 \leq (i-4) + j + (n-i-j-2)$, i.e., $d_2 \leq n-3$.

(iii) if $d_1 = i+j = n-2$ and $i \geq 6$, then $\pi^* = (d_1-5, d_2-3, 3^{i-4}, 2^{n-i-2})$. Therefore, $d_2 - 3 \leq (i-4) + (n-i-2)$, i.e. $d_2 \leq n-3$;

(6) If $\pi = ((n-1)^2, 4, 3^{n-3})(n \geq 7 \ and \ n \ is \ odd)$ is potentially $K_6 - C_5$-graphic, then $\pi^* = (n-6, n-4, 1, 3^{n-6})$ is graphic, hence $n-4-1 \leq n-6$, a contradiction.

(7) If $\pi = (n-1, 3^6, 1^{n-7})(n \geq 7)$ is potentially $K_6 - C_5$-graphic, then $\pi^* = (n-6, 3, 1^{n-7})$ is graphic, hence $\pi_1 = (2) \in GS_n$, a contradiction.

If $\pi = (n-1, 3^7, 1^{n-8})(n \geq 8)$is potentially $K_6 - C_5$-graphic, then $\pi^* = (n-6, 3^2, 1^{n-8})$ is graphic, hence $\pi_1 = (2^2) \in GS_n$, a contradiction.

If $\pi =$$(5,4,3^5)$,$(5^2,3^5,1)$,$(5,4^2,3^5)$,$(5,4,3^7)$,$(6,3^7,1)$,$(5^2,3^6)$,$(5^2,4,3^4)$, $(5,3^7)$,$(6,4,3^6)$,$(5^2,3^4,2)$,$(5,3^6,1)$,$(5,3^6,2,1)$,$(5,3^7,2)$,$(5,3^8,1)$,$(5,4,3^5,2)$, $(5,4,3^6,1)$,$(5,3^5,2)$,$(5,3^5,2^2)$,$(6,5,3^5)$,$(5,3^7,1^2)$,$(6,3^8)$, $(6^2,3^4,2)$, then it is easy to see that they are not potentially $K_6 - C_5$-graphic .

Now we prove the sufficient condition. Suppose the graphic sequence π satisfies the conditions (1) to (7). Our proof is by induction on n.

We first prove the base case where $n = 6$. Then π is one of the following: (5^6),$(5^4,4^2)$,$(5^3,4^2,3)$,$(5^3,3^3)$,$(5^2,4^4)$,$(5^2,4^2,3^2)$, $(5^2,3^4)$,$(5,4^4,3)$, $(5,4^2,3^3)$, $(5,3^5)$. It is easy to check that others are potentially $K_6 - C_5$-graphic except for $(5^3,3^3)$ and $(5^2,3^4)$. Now suppose that the sufficiency holds for $n-1 (n \geq 7)$, we will show that π is potentially $K_6 - C_5$-graphic in terms of the following cases:

Case 1: $d_n \geq 6$

Consider $\pi' = (d'_1, d'_2, \cdots, d'_{n-1})$ where $d'_{n-1} \geq 5$.

It is easy to check that π' satisfies (1) to (7). Then by the induction hypothesis, π' is potentially $K_6 - C_5$-graphic, and hence so is π.

Case 2: $d_n = 5$

Consider $\pi' = (d'_1, d'_2, \cdots, d'_{n-1})$ where $d'_1 \geq 5$ and $d'_{n-1} \geq 4$.

It is easy to check that π' satisfies (1) to (7). Then by the induction hypothesis, π' is potentially $K_6 - C_5$-graphic, and hence so is π.

Case 3: $d_n = 4$

Consider $\pi' = (d'_1, d'_2, \cdots, d'_{n-1})$ where $d'_1 \geq 4$ and $d'_{n-1} \geq 3$.

It is easy to check that π' satisfies (2) to (7). If π' satisfies (1), then by the induction hypothesis, π' is potentially $K_6 - C_5$-graphic, and hence so is π.

If π' does not satisfy (1), then $d'_1 = 4$. Therefore, $d_5 = 4 = d_6 = \cdots = d_{n-1}$. Then $\pi =(5^2, 4^{n-2})(n \geq 7)$ or $\pi =(5^4, 4^{n-4})(n \geq 7)$. According to Lemma 2.8 and 2.7, $\pi =(5^2, 4^{n-2})(n \geq 7)$ and $\pi =(5^4, 4^{n-4})(n \geq 7)$ holding the conditions of Theorem 3.1 are potentially $K_6 - C_5$-graphic.

Case 4: $d_n = 3$

Consider $\pi' = (d'_1, d'_2, \cdots, d'_{n-1})$ where $d'_1 \geq 4$, $d'_{n-3} \geq 3$ and $d'_{n-1} \geq 2$.

It is easy to check that π' satisfies (2). If π' satisfies (1),(3) to (7), then by the induction hypothesis, π' is potentially $K_6 - C_5$-graphic, and hence so is π.

If π' does not satisfy (1), then $d'_1 = 4$ or $d'_6 \leq 2$. If $d'_1 = 4$, then $d_1 = 5$ and $3 \leq d_4 \leq 4$. Therefore, $\pi =(5^i, 4^j, 3^{n-i-j})(1 \leq i \leq 3, j \geq 0, n \geq 7)$.

According to Lemma 2.7 to 2.9, $\pi =(5^i, 4^j, 3^{n-i-j})(1 \leq i \leq 3, j \geq 0, n \geq 7)$ holding the conditions of Theorem 3.1 is potentially $K_5 - C_4$-graphic.

If $d'_6 = 2$, then $d_3 = 3 = \cdots = d_n$ .Therefore, $\pi =(d_1, d_2, 3^{n-2})(d_1 \geq 5, d_2 \geq 3, n \geq 7)$. According to Lemma 2.10, $\pi =(d_1, d_2, 3^{n-2})(d_1 \geq 5, d_2 \geq 3)$ holding the conditions of Theorem 3.1 is potentially $K_6 - C_5$-graphic.

If π' does not satisfy (3), then $n = 7$ and $\pi' = (5^2, 3^4)$, hence $\pi =(6, 6, 4, 3^4)$, a contradiction to the condition (6).

If π' does not satisfy (4), then π' satisfies (4)(ii), (4)(iii) and (4)(iv).

If π' does not satisfy (4)(i), then $\pi' = ((n-1)^2, d'_3, 3^{n-3})$ where $i \geq 4$ and $d'_3 \geq 5$, hence $\pi = (n^2, d'_3 + 1, 3^{n-2})$, a contradiction to the condition (4)(i).

If π' does not satisfy (5), then π' satisfies (5)(i).

If π' does not satisfy (5)(ii), then $\pi' = (n-1, n-2, 3^{n-j-2}, 2^j)$ or $\pi' = ((n-1)^2, 3^{n-j-2}, 2^j)$ where $n-j-2 \geq 5$ and $j \leq 1$. If $\pi' = (n-1, n-2, 3^{n-2})$, then $\pi = (n, n-1, 4, 3^{n-2})(n \geq 7$ and n is odd) is potentially $K_6 - C_5$-graphic . If $\pi' = (n-1, n-2, 3^{n-3}, 2)$, then $\pi = (n, n-1, 3^{n-1})$, a contradiction to the condition (5)(i). If $\pi' = ((n-1)^2, 3^{n-2})$, then $\pi = (n^2, 4, 3^{n-2})$, a contradiction to the condition (6). If $\pi' = ((n-1)^2, 3^{n-3}, 2)$, then $\pi = (n^2, 3^{n-1})$, a contradiction to the condition (5)(ii).

If π' does not satisfy (5)(iii), then $\pi' = ((n-2)^2, 3^{n-2})(n \geq 8)$ or $\pi' = ((n-2)^2, 3^{n-3}, 2)(n \geq 9)$. Therefore, $\pi = ((n-1)^2, 4, 3^{n-2})(n \geq 8)$ is potentially K_6-C_5-graphic or $\pi=((n-1)^2,3^{n-1})(n\geq 9)$ which contradicts the condition (5)(*iii*).

If π' does not satisfy (6), then $\pi=(n^2,5,3^{n-2})$ which contradicts the condition (4)(*i*) or $\pi=(n^2,4^2,3^{n-3})$ is potentially K_6-C_5-graphic .

If π' does not satisfy (7), then $\pi'=(6,3^6)$,$(7,3^7)$,$(5,4,3^5)$,$(5,4^2,3^5)$, $(5,4,3^7)$, $(5^2,3^6)$,$(5^2,4,3^4)$,$(5,3^7)$,$(6,4,3^6)$,$(5^2,3^4,2)$,$(5,3^7,2)$,$(5,4,3^5,2)$, $(5,3^5,2)$, $(5,3^5,2^2)$, $(6,5,3^5)$, $(6,3^8)$, $(6^2,3^4,2)$, hence $\pi=(7,4^2,3^5)$, $(8,4^2,3^6)$, $(6,5,4,3^5)$,$(6,4^3,3^4)$,$(6,5^2,3^6)$,$(6,5,4^2,3^5)$,$(6,4^4,3^4)$,$(6,5,4,3^7)$, $(6,4^3,3^6)$, $(6^2,4,3^6)$, $(6^2,5,3^5)$, $(6^2,4^2,3^4)$, $(6,4^2,3^6)$, $(7,5,4,3^6)$, $(7,4^3,3^5)$, $(6^2,3^6)$, $(6,4,3^8)$, $(6,5,3^7)$, $(6,4,3^6)$,$(6,3^8)$,$(7,6,4,3^5)$,$(7,4^2,3^7)$,$(7^2,4,3^5)$, $(7^2,3^6)$. It is easy to check that others are potentially K_6-C_5-graphic except $(6^2,3^6)$, $(6,4,3^6)$, $(6,3^8)$, $(7^2,4,3^5)$ and $(7^2,3^6)$. Then $\pi=(6^2,3^6)$,$(6,4,3^6)$ $(6,3^8)$,$(7^2,4,3^5)$ and $(7^2,3^6)$, a contradiction to the conditions (5)(*iii*), (7), (4)(*iv*), (6) and (5)(*ii*).

Case 5: $d_n=2$

Consider $\pi'=(d'_1,d'_2,\cdots,d'_{n-1})$ where $d'_1\geq 4$, $d'_5\geq 3$ and $d'_{n-1}\geq 2$.

It is easy to check that π' satisfies (2) and (3). If π' satisfies (1) and (4) to (7), then by the induction hypothesis, π' is potentially K_6-C_5-graphic, hence so is π.

If π' does not satisfy (1), then $d'_1=4$ or $d'_6\leq 2$. If $d'_1=4$, then $d_1=5$ and $d_3\leq 4$. Therefore, $\pi=(5^2,4^i,3^j,2^{n-2-i-j})(i+j\geq 4,n\geq 7)$ or $\pi=(5,4^i,3^j,2^{n-1-i-j})(i+j\geq 5,n\geq 7)$.

According to Lemma 2.8 and 2.9, $\pi=(5^2,4^i,3^j,2^{n-2-i-j})(i+j\geq 4,n\geq 7)$ and $\pi=(5,4^i,3^j,2^{n-1-i-j})(i+j\geq 5,n\geq 7)$ holding the conditions of Theorem 3.1 are potentially K_6-C_5-graphic.

If $d'_6=2$, then $d_2=3=d_3=d_4=d_5=d_6$ and $d_7=\cdots=d_n$. Therefore, $\pi=(d_1,3^5,2^{n-6})(d_1\geq 5,n\geq 7)$.

According to Lemma 2.11, $\pi=(d_1,3^5,2^{n-6})(d_1\geq 5,n\geq 7)$ holding the conditions of Theorem 3.1 is potentially K_6-C_5-graphic.

If π' does not satisfy (4), then π' satisfies (4)(*ii*) and (4)(*iii*).

If π' does not satisfy (4)(*i*), then $\pi'=((n-1)^2,d'_3,3^{n-3})$ where $i\geq 4$ and $d'_3\geq 5$. If $d'_3\neq n-2$, then $\pi=(n^2,d'_3,3^{n-3},2)$, a contradiction to the condition (4)(i). If $d'_3=n-2$, then $\pi=(n^2,n-2,3^{n-3},2)$ which contradicts the condition (2) or $\pi=(n,(n-1)^2,3^{n-3},2)$ which also contradicts the condition (2).

If π' does not satisfy (4)(*iv*), then $\pi'=((n-1)^2,d'_3,3^i,2^{n-i-3})$ where $i\geq 5$, $d'_3\geq 4$ and $d'_2\geq d'_3+2$. Therefore, $\pi=(n^2,d'_3,3^i,2^{n-i-2})$, a contradiction to the condition (4)(iv).

If π' does not satisfy (5), then π' satisfies (5)(i).

If π' does not satisfy (5)(ii) where $i \geq 4$ and $d'_2 \geq 5$, then $\pi' = (d'_1, d'_2, 3^i, 2^{n-i-2})$. Therefore, $\pi = (d'_1+1, d'_2+1, 3^i, 2^{n-i-1})$, a contradiction to the condition (5)(ii).

If π' does not satisfy (5)(iii), then $\pi' = ((n-2)^2, 3^i, 2^{n-i-2})(i \geq 6)$. Therefore, $\pi = ((n-1)^2, 3^i, 2^{n-i-2+1})(i \geq 6)$, a contradiction to the condition (5)(iii).

If π' does not satisfy (6), then $\pi =(n^2, 4, 3^{n-3}, 2)$, a contradiction to the condition (4)(iii).

If π' does not satisfy (7), then $\pi' = (6, 3^6)$,$(7, 3^7)$,$(5, 4, 3^5)$, $(5, 4^2, 3^5)$, $(5, 4, 3^7)$, $(5^2, 3^6)$,$(5^2, 4, 3^4)$,$(5, 3^7)$, $(6, 4, 3^6)$,$(5^2, 3^4, 2)$, $(5, 3^7, 2)$,$(5, 4, 3^5, 2)$, $(5, 3^5, 2)$, $(5, 3^5, 2^2)$, $(6, 5, 3^5)$, $(6, 3^8)$, $(6^2, 3^4, 2)$, hence $\pi =(7, 4, 3^5, 2)$, $(8, 4, 3^6, 2)$, $(6, 4^2, 3^4, 2)$, $(6, 5, 3^5, 2)$, $(6, 5, 4, 3^5, 2)$, $(5^3, 3^5, 2)$, $(6, 4^3, 3^4, 2)$, $(6, 5, 3^7, 2)$, $(6, 4^2, 3^6, 2)$, $(6^2, 3^6, 2)$, $(6^2, 4, 3^4, 2)$, $(6, 5^2, 3^4, 2)$, $(6, 4, 3^6, 2)$, $(7, 5, 3^6, 2)$, $(7, 4^2, 3^5, 2)$, $(6^2, 3^4, 2^2)$, $(6, 4, 3^6, 2^2)$, $(6, 5, 3^5, 2^2)$, $(6, 4^2, 3^4, 2^2)$, $(6, 4, 3^4, 2^2)$, $(6, 4, 3^4, 2^3)$, $(7, 6, 3^5, 2)$,$(7, 4, 3^7, 2)$,$(7^2, 3^4, 2^2)$. It is easy to check that others are potentially K_6-C_5-graphic except $(6^2, 3^4, 2^2)$, $(7, 6, 3^5, 2)$ and $(7^2, 3^4, 2^2)$. Then $\pi =(6^2, 3^4, 2^2)$,$(7, 6, 3^5, 2)$ and $(7^2, 3^4, 2^2)$, a contradiction to the conditions (5)(iii) and (5)(ii).

Case 6: $d_n = 1$

Consider $\pi' = (d'_1, d'_2, \cdots, d'_{n-1})$ where $d'_1 \geq 4$, $d'_6 \geq 3$ and $d'_{n-1} \geq 1$.

It is easy to check that π' satisfies (2) and (3). If π' satisfies (1) and (4) to (7), then by the induction hypothesis, π' is potentially $K_6 - C_5$-graphic, and hence so is π.

If π' does not satisfy (1), then $d'_1 = 4$. If $d'_1 = 4$, then $d_1 = 5$. Therefore, $\pi =(5, 4^i, 3^j, 2^k, 1^{n-i-j-k-1})(i+j \geq 5, n \geq 7)$.

According to Lemma 2.9, $\pi =(5, 4^i, 3^j, 2^k, 1^{n-i-j-k-1})(i+j \geq 5, n \geq 7)$ holding the conditions of Theorem 3.1 is potentially $K_6 - C_5$-graphic.

If π' does not satisfy (4)(i), then $\pi' = ((n-1)^2, d'_3, 3^{n-3})$ where $i \geq 4$ and $d'_3 \geq 5$.

If $d'_3 \neq n-2$, then $\pi = (n, n-1, d'_3, 3^{n-3}, 1)$, a contradiction to the condition (4)(iii). If $d'_3 = n-2$, then $\pi = (n, n-1, n-2, 3^{n-3}, 1)$ or $\pi = ((n-1)^3, 3^{n-3}, 1)$ which also contradicts the conditions (4)(iii) and (2).

If π' does not satisfy (4)(ii), then $\pi' = (d'_1, d'_2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$ where $i \geq 4$, $d'_3 \geq 4$ and $d'_2 \geq d'_3 + 2$.

If $d'_1 \neq d'_2 + 1$, then $\pi = (d'_1 + 1, d'_2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$, a contradiction to the condition $(4)(ii)$. If $d'_1 = d'_2 + 1$, then $\pi = (d'_1 + 1, d'_2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$ which contradicts the condition $(4)(ii)$ or $\pi = (d'_1, d'_2 + 1, d'_3, 3^i, 2^j, 1^{n-i-j-3})$ which also contradicts the condition $(4)(ii)$.

If π' does not satisfy $(4)(iii)$, then $\pi' = (d'_1, d'_2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$ and $i \geq 4$, $d'_3 \geq 5$, $d'_2 \geq d'_3 + 2$.

If $d'_1 \neq d'_2 + 1$, then $\pi = (d'_1 + 1, d'_2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$, which contradicts the condition $(4)(iii)$. If $d'_1 = d'_2 + 1$, then $\pi = (d'_1 + 1, d'_2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$ or $\pi = (d'_1, d'_2 + 1, d'_3, 3^i, 2^j, 1^{n-i-j-3})$, which contradicts the condition $(4)(iii)$.

If π' does not satisfy $(4)(iv)$, then $\pi' = ((n-1)^2, d'_3, 3^i, 2^j, 1^{n-i-j-3})$ and $i \geq 4$, $d'_2 \geq d'_3 + 2$. Then $\pi = (n, n-1, d'_3, 3^i, 2^j, 1^{n-i-j-2})$, which contradicts the conditions $(4)(ii)$ and $(4)(iii)$.

If π' does not satisfy $(5)(i)$, then $\pi' = (d'_1, d'_2, 3^i, 2^j, 1^{n-i-j-2})$ where $i \geq 5$ and $d'_2 \geq 5$. If $d'_1 \neq d'_2 - 1$, then $\pi = (d'_1 + 1, d'_2, 3^i, 2^j, 1^{n-i-j-1})$, a contradiction to the condition $(5)(i)$. If $d'_1 = d'_2 - 1$, then $\pi = (d'_1 + 1, d'_2, 3^i, 2^j, 1^{n-i-j-1})$ which contradicts the condition $(5)(i)$ or $\pi = (d'_1, d'_2 + 1, 3^i, 2^j, 1^{n-i-j-1})$ which also contradicts the condition $(5)(i)$.

If π' does not satisfy $(5)(ii)$, then $\pi' = (n-1, n-2, 3^i, 2^{n-i-2})$,$((n-1)^2, 3^i, 2^{n-i-2})$ or $((n-2)^2, 3^i, 2^{n-i-3}, 1)$. If $\pi' = (n-1, n-2, 3^i, 2^{n-i-2})$, then $\pi = (n, n-2, 3^i, 2^{n-i-2}, 1)$ which contradicts the condition $(5)(i)$ or $\pi = ((n-1)^2, 3^i, 2^{n-i-2}, 1)$ which contradicts the condition $(5)(ii)$. If $\pi' = ((n-1)^2, 3^i, 2^{n-i-2})$, then $\pi = (n, n-1, 3^i, 2^{n-i-2}, 1)$, a contradiction to the condition $(5)(i)$. If $\pi' = ((n-2)^2, 3^i, 2^{n-i-3}, 1)$, then $\pi = (n-1, n-2, 3^i, 2^{n-i-2}, 1^2)$, a contradiction to the condition $(5)(i)$.

If π' does not satisfy $(5)(iii)$, then $\pi' = ((n-2)^2, 3^i, 2^{n-i-2}) (i \geq 6)$. Therefore, $\pi = (n-1, n-2, 3^i, 2^{n-i-2}, 1)$ $(i \geq 6)$ is potentially $K_6 - C_5$-graphic.

If π' does not satisfy (6), then $\pi =(n, n-1, 4, 3^{n-3}, 1)$, a contradiction to the condition $(4)(ii)$.

If π' does not satisfy (7), then $\pi' =(n-1, 3^6, 1^{n-7})(n \geq 7)$,$(n-1, 3^7, 1^{n-8})(n \geq 8)$, $(5, 4, 3^5)$, $(5^2, 3^5, 1)$, $(5, 4^2, 3^5)$, $(5, 4, 3^7)$, $(6, 3^7, 1)$, $(5^2, 3^6)$, $(5^2, 4, 3^4)$, $(5, 3^7)$,$(6, 4, 3^6)$, $(5^2, 3^4, 2)$,$(5, 3^6, 1)$,$(5, 3^6, 2, 1)$,$(5, 3^7, 2)$,$(5, 3^8, 1)$, $(5, 4, 3^5, 2)$, $(5, 4, 3^6, 1)$, $(5, 3^5, 2)$, $(5, 3^5, 2^2)$, $(6, 5, 3^5)$, $(5, 3^7, 1^2)$, $(6, 3^8)$, $(6^2, 3^4, 2)$. Hence $\pi =(n, 3^6, 1^{n-6})(n \geq 7)$,$(n, 3^7, 1^{n-7})(n \geq 8)$, $(6, 4, 3^5, 1)$, $(5^2, 3^5, 1)$, $(6, 5, 3^5, 1^2)$, $(6, 4^2, 3^5, 1)$, $(5^2, 4, 3^5, 1)$, $(6, 4, 3^7, 1)$, $(5^2, 3^7, 1)$, $(7, 3^7, 1^2)$, $(6, 5, 3^6, 1)$, $(6, 5, 4, 3^4, 1)$, $(5^3, 3^4, 1)$, $(6, 3^7, 1)$, $(7, 4, 3^6, 1)$, $(6, 5, 3^4, 2,$

$1)$, $(6,3^6,1^2)$, $(6,3^6,2,1^2)$, $(6,3^7,2,1)$, $(6,3^8,1^2)$, $(6,4,3^5,2,1)$, $(5^2,3^5,2,1)$, $(6,4,3^6,1^2)$, $(5^2,3^6,1^2)$, $(6,3^5,2,1)$, $(6,3^5,2^2,1)$, $(7,5,3^5,1)$, $(6^2,3^5,1)$, $(6,3^7,1^3)$, $(7,3^8,1)$,$(7,6,3^4,2,1)$. It is easy to check that others are potentially K_6-C_5-graphic except $\pi=(n,3^6,1^{n-6})(n\geq 7)$, $(n,3^7,1^{n-7})(n\geq 8)$, $(5^2,3^5,1)$, $(6,3^7,1)$, $(7,5,3^5,1)$, $(6^2,3^5,1)$ and $(7,6,3^4,2,1)$. Then $\pi=(n,3^6,1^{n-6})(n\geq 7)$, $(n,3^7,1^{n-7})(n\geq 8)$, $(5^2,3^5,1)$, $(6,3^7,1)$, $(7,5,3^5,1)$, $(6^2,3^5,1)$ and $(7,6,3^4,2,1)$, a contradiction to the conditions (7), (5) and (4).

Corollary 3.2 For $n\geq 6$, $\sigma(K_6-C_5,n)=6n-10$.

Proof: First we claim $\sigma(K_6-C_5,n)\geq 6n-10$ for $n\geq 6$. We would like to show there exists π_1 with $\sigma(\pi_1)=6n-12$ such that π_1 is not potentially K_6-C_5-graphic. Let $\pi_1=((n-1)^3,3^{n-3})$. It is easy to see that $\sigma(\pi_1)=6n-12$ and π_1 is not potentially K_6-C_5-graphic by Theorem 3.1.

Now we easily show if π is an n-term $(n\geq 6)$ graphic sequence with $\sigma(\pi)\geq 6n-10$, then there exists a realization of π containing a K_6-C_5.

(1) We claim that $d_1\geq 5$ and $d_6\geq 3$. By way of contradiction, we assume that $d_1\leq 4$ or $d_6\leq 2$. If $d_1\leq 4$, then $\sigma(\pi_1)\leq 4\times(n-1)< 6n-12(n\geq 6)$, a contradiction. If $d_6\leq 2$, then $\sigma(\pi_1)=\sum_{i=1}^{5}d_i+\sum_{i=6}^{n}d_i\leq 5\times(5-1)+\sum_{i=6}^{n}min(5,d_i)+\sum_{i=6}^{n}d_i=20+2\sum_{i=6}^{n}d_i\leq 4\text{n}<6n-10(n\geq 6)$, a contradiction.

(2) If $\pi=(d_1,d_2,d_3,3^i,2^j,1^{n-i-j-3})$ where $i\geq 3$ and $d_3\geq 5$, then $\sigma(\pi)=d_1+d_2+d_3+3i+2j+n-i-j-3\leq 3\times(n-1)+3\times(n-3)=6n-12$, a contradiction. Then π with $\sigma(\pi)\geq 6n-10$ satisfies the condition (2) of the Theorem 3.1.

Similar to prove that π with $\sigma(\pi)\geq 6n-10$ satisfies the conditions (3) to (7) of the Theorem 3.1. According to the Theorem 3.1, π is potentially K_6-C_5-graphic.

Acknowledgment

The authors wish to thank Professor Gang Chen, R.J. Gould, Jiongsheng Li, Rong Luo, John R. Schmitt, Zi-Xia Song, Amitabha Tripathi, Jianhua Yin and Mengxiao Yin for sending some their papers to us.

References

[1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, The Macmillan Press Ltd., 1976.

[2] P.Erdös, M.S. Jacobson and J. Lehel, Graphs realizing the same degree sequences and their respective clique numbers, in Graph Theory, Combinatorics and Application, Vol. 1(Y. Alavi et al., eds.), John Wiley and Sons, Inc., New York, 1991, 439-449.

[3] R.J. Gould, M.S. Jacobson and J. Lehel, Potentially G-graphic degree sequences,in Combinatorics, Graph Theory and Algorithms,Vol. 2 (Y. Alavi et al.,eds.), New Issues Press, Kalamazoo, MI, 1999, 451-460.

[4] Lili Hu and Chunhui Lai , on potentially $K_5 - C_4$-graphic sequences, accepted by Ars Combinatoria.

[5] Lili Hu and Chunhui Lai , on potentially $K_5 - E_3$-graphic sequences, accepted by Ars Combinatoria.

[6] Lili Hu, Chunhui Lai and Ping Wang, on potentially $K_5 - H$-graphic sequences, accepted by Czechoslovak Mathematical Journal.

[7] Lili Hu and Chunhui Lai , on potentially 3-regular graph graphic sequences, accepted by Utilitas Mathematica.

[8] D.J. Kleitman and D.L. Wang , Algorithm for constructing graphs and digraphs with given valences and factors,Discrete Math., 6(1973),79-88.

[9] Chunhui Lai, A note on potentially $K_4 - e$ graphical sequences,Australasian J. of Combinatorics 24(2001),123-127.

[10] Chunhui Lai, An extremal problem on potentially $K_m - C_4$-graphic sequences, Journal of Combinatorial Mathematics and Combinatorial Computing, 61 (2007), 59-63.

[11] Chunhui Lai, An extremal problem on potentially $K_m - P_k$-graphic sequences, accepted by International Journal of Pure and Applied Mathematics.

[12] Chunhui Lai, An extremal problem on potentially $K_{p,1,1}$-graphic sequences, Discrete Mathematics and Theoretical Computer Science 7(2005), 75-81.

[13] Chunhui Lai and Lili Hu, An extremal problem on potentially $K_{r+1}-H$-graphic sequences, accepted by Ars Combinatoria.

[14] Chunhui Lai, The smallest degree sum that yields potentially $K_{r+1}-Z$-graphical Sequences, accepted by Ars Combinatoria.

[15] Jiong-Sheng Li and Zi-Xia Song, The smallest degree sum that yields potentially P_k-graphical sequences, J. Graph Theory, 29(1998), 63-72.

[16] Jiong-sheng Li, Zi-Xia Song and Rong Luo, The Erdös-Jacobson-Lehel conjecture on potentially P_k-graphic sequence is true, Science in China(Series A), 41(5)(1998), 510-520.

[17] Jiong-Sheng Li and Zi-Xia Song, on the potentially P_k-graphic sequences, Discrete Math., 195(1999), 255-262.

[18] Jiong-Sheng Li and Zi-Xia Song, An extremal problem on the potentially P_k-graphic sequences, Discrete Math., 212(2000), 223-231.

[19] Jiong-sheng Li and Jianhua Yin, A variation of an extremal theorem due to Woodall, Southeast Asian Bulletin of Math., 25(2001), 427-434.

[20] Rong Luo, On potentially C_k-graphic sequences, Ars Combinatoria 64(2002), 301-318.

[21] Rong Luo, Morgan Warner, On potentially K_k-graphic sequences, Ars Combin. 75(2005), 233-239.

[22] Elaine M. Eschen and Jianbing Niu, On potentially K_4-e-graphic sequences, Australasian Journal of Combinatorics, 29(2004), 59-65.

[23] Zhenghua Xu,Potentially H graphic sequences,Journal of Zhangzhou Normal University(Natural Science),21(3)(2008),7-9.

[24] Zhenghua Xu, Characterizations On Potentially K_6-H-graphic Sequences, A Dissertation for the Degree of M.Science of Zhangzhou Normal University, 2009,10.

[25] J.H.Yin, The Smallest Degree Sum Yielding Potentially $K_{1,1,3}$-graphic Positive Sequences, Natural Science Journal of Hainan University, 22(2004),200-204.

[26] Jianhua Yin and Jiongsheng Li, Two sufficient conditions for a graphic sequence to have a realization with prescribed clique size, Discrete Math.,301(2005) 218-227.

[27] Jianhua Yin, Jiongsheng Li and Rui Mao, An extremal problem on the potentially $K_{r+1} - e$-graphic sequences, Ars Combinatoria 74(2005), 151-159.

[28] J.H.Yin and J.S.Li, The smallest degree sum that yields potentially $K_{r,r}$-graphic sequences, Sci. China Ser. A, 45(2002), 694-705.

[29] J.H.Yin and J.S.Li, An extremal problem on the potentially $K_{r,s}$-graphic sequences, Discrete Math., 26(2003), 295-305.

[30] Jianhua Yin and Gang Chen, On potentially $K_{r_1,r_2,\cdots,r_m}$-graphic sequences, Utilitas Mathematica, 72(2007), 149-161.

[31] Mengxiao Yin, The smallest degree sum that yields potentially $K_{r+1} - K_3$-graphic sequences, Acta Math. Appl. Sin. Engl. Ser. 22(2006), no. 3, 451-456.

[32] Mengxiao Yin, Jianhua Yin, On potentially H-graphic sequences, Czechoslovak Mathematical Journal, 57(2)(2007),705-724.

[33] M.X.Yin, A characterization on potentially $K_6 - E(K_3)$-graphic sequences, accepted by Ars Combinatoria.